

Ramkhamhaeng (pronounced ram-cam-hang) Hospital sits at the center of one of Thailand's largest and most respected hospital groups. With 350 beds and eight sister hospitals Ramkhamhaeng has the resources, experience and expertise to provide international quality health care at very competitive prices. This combined with a long tradition of Thai hospitality has earned Ramkhamhaeng Hospital the confidence of the people of Thailand, the Region, and the World.

Thailand has gained renown for all forms of cosmetic surgery, cardiac care, and other types of specialized medicine. Ramkhamhaeng Hospital has many internationally trained doctors with world-class credentials. In this era of jet travel, computers, and the Internet, Globalized medicine has arrived. We offer world class healthcare in a beautiful tropical setting at a fraction of the cost of the USA or Europe. If you pay your own medical bills because it is an elective surgery or you are uninsured and don't have or can't get insurance

[Contact Us.](#)



WWW.RamHospital.com/Index.html

1238 Ramkhamhaeng Road
Huamark, Bangkok, Bangkok
Thailand 10240

Email: The_DocBangkok@yahoo.com



**The Smart
Choice For
Global
Healthcare**

**The Healthcare You Need At
Prices You Can Afford**

VISITING HOURS

Ramkhamhaeng Hospital recognizes that family and friends are important to support in your recovery. Visiting hours are flexible and are based on your needs. An important aspect of recovery is to be sure you are receiving adequate rest. You have the right to limit visitors at any time.

It is recommended that visitors other than immediate family support limit their stay to 15 minutes with a maximum of two visitors at a time. Children may visit if they are free of illness and supervised by an adult.

WWW.RamHospital.com/Index.html

1238 Ramkhamhaeng Road
Huamark, Bangkok, Bangkok
Thailand 10240

Email: The_DocBangkok@yahoo.com

THE SMART CHOICE FOR
GLOBAL HEALTHCARE



Gender Reassignment Clinic
The Healthcare You Need At
Prices You Can Afford

Our price for [Gender Reassignment Surgery](#) is \$6250 USD. This compares to up to \$40,000 in the USA.

WWW.RamHospital.com/Index.html

1238 Ramkhamhaeng Road
Huamark, Bangkok, Bangkok
Thailand 10240

Email: The_DocBangkok@yahoo.com





Gender Reassignment Clinic



Male to Female (MtoF)

Transsexuals are people who have a long-standing and persistent feeling that one's sexual identity is inconsistent with one's anatomic sex.

You must be on hormone therapy for 6 months before surgery.

Male to female transsexuals use the hormone **estrogen** to feminize their voices, faces, and body characteristics. This hormone changes their fat distribution, increases their breast size, and decreases their body hair. One thing that this hormone does not do, however, is to eliminate facial hair. Estrogen also shrinks the penis and testicles. Any medical doctor, including a psychiatrist, can prescribe hormones once they determine that their patient is suitable for them.

A vaginoplasty involves the removal of the testes to create labia. It also involves the inversion of the penis to create a vagina -- this requires extensive surgery and follow-up visits. The good news is that these newly constructed vaginas look amazingly like the real thing.

Patients who are suitable for Gender Reassignment must meet the following specifications:

1. Must be HIV NEGATIVE
2. Have been living for at least one year full-time in the new gender role. Living in this role should be complete and successful.
3. Live with desire for being the other gender for more than 2 years.
4. Take hormone therapy for at least 6 months.
5. Gain a recommendation from a psychiatrist or therapist.
6. Have a negative attitude of your current sexual organs.
7. No psychiatric illness.

Preparation before operation :

1. Should stop taking hormone pill for 2 weeks before operation.
2. Stop smoking and taking Aspirin because this will cut down on the amount of bleeding during the procedure.
3. The patient should eat a soft diet for 2-3 days prior to surgery to prevent the problem of defecation after operation.
4. The patient should not drink and eat food for 6 hours before the operation.



Penile Inversion Vaginoplasty

This is widely regarded as the preferred method of male-to-female GRS, is widely performed, and when performed by an accomplished surgeon, can produce very satisfactory results.

The basic method involves turning the penile skin 'inside out' and using it to line a vaginal cavity created by blunt dissection through the muscles of the perineal area. The penis and testes are removed. Many surgeons use variants of the technique: a pure penile inversion limits the size of the vagina that can be created, depending on the amount of penile skin available. In many patients it is necessary to supplement this material with scrotal skin or by means of a skin graft, often taken from the thigh or abdomen.

A modern method involves the use of scrotal tissue, and is sometimes termed 'peno-scrotal inversion'. This is widely regarded as the preferred method. In this technique, the penile skin is usually divided, part being used to form the floor of the vulva, and part being used to form the anterior wall of the vagina, with a flap of scrotal tissue being used to form the posterior wall and apex of the vagina. This has the considerable advantage that a satisfactory vagina can usually be formed even in circumcised or mildly-intersexed patients where penile tissue may be very limited; the corresponding disadvantage is that the risk of vaginal prolapse increases with the proportion of scrotal tissue used, as scrotal tissue adheres less well to the muscles than penile tissue.

Labia are constructed from scrotal tissue, and the urethra is shortened and everted to correspond to normal female anatomy. A good surgeon will re-site the urethral meatus to an appropriate female position, and will ensure that the erectile material of the penis is removed to the maximum possible extent, so as not to leave an unsightly and possibly painful 'stump'. The remnant of the prostate gland (long-term hormone therapy causes it to atrophy and shrink) is left in place, and provides sexual sensation through the anterior wall of the vagina, analogous to the so-called 'G spot' of natural-born females.

After this operation, the patient will have a pack (usually of surgical gauze) placed in the vagina, to retain the tissues in their proper locations for about five days while healing proceeds. During this time she will be on bed rest, and a clear-fluid-only diet to avoid the possibility of a bowel movement which could damage the vagina or disturb the pack. The patient will also be catheterised, typically for seven days, to allow the urethra to heal.

Many male to female transsexuals also have cosmetic surgery to feminize their facial features and reduce the size of their Adam's apple. Other surgeries can include: suction-assisted lipoplasty of the waist, rhinoplasty, facial bone reduction, face-lift, and blephroplasty. Ramkham-

WWW.RamHospital.com/Index.html

1238 Ramkhamhaeng Road
Huamark, Bangkok
Thailand 10240

Email: The_DocBangkok@Yahoo.com

